

## JOSEPH D. LAMPLOT, M.D. ORTHOPAEDIC SURGERY & SPORTS MEDICINE

Appointments: (847) 238-2812 | SPORTSMEDICINE@NORTHSHORE.ORG

21481 N. RAND RD, KILDEER, IL 60047 1535 ELLINWOOD AVE, DES PLAINES, IL 60016 CLINICAL CONCERNS: (331) 221-6385

Date of Surgery:

## Patellar or Quadriceps Tendon Reconstruction

<u>Postop</u>	<u>Goals</u>	<u>Precautions</u>	<u>Exercises</u>
Weeks 0-2 PT 1-2x/week HEP daily	Edema and pain control Protect surgical repair Maintain full extension Limit quad inhibition	TTWB Brace locked in extension at all times for ambulation and during sleep - remove for exercises only No active or passive knee ROM Avoid prolonged standing/walking Ensure proper brace alignment	Brace education Cryotherapy Proximal/distal strengthening Quadriceps re-education & isometrics SLR brace locked in extension Scar mobilization Patella mobilization Gait training
Weeks 2-6 PT 1-2x/week HEP daily	Edema and pain control Protect surgical repair Maintain full extension Limit quad inhibition Goal: ROM 0-60°	PWB (50%), progress to WBAT Brace locked in extension at all times for ambulation and during sleep - remove for exercises only No active knee extension Avoid prolonged standing/walking Ensure proper brace alignment  Must adhere to MD ROM limit Knee flexion progression: 0-30° starting in week 3 Goal of 60° by 6 weeks	Continue Phase 1 exercises Seated A/AA knee FLEXION within limits Passive knee EXTENSION Emphasize adherence to ROM limits, brace compliance
Weeks 6-9 PT 2-3x/week HEP daily	Goal: ROM 0-90° No extensor lag Normalize gait Wean from gait aids Minimize pain and swelling	Brace locked in extension at all times for ambulation and during sleep - remove for exercises only  Notify MD if knee flexion <90° by 9 weeks	Continue Phase 2 exercises Active flexion/passive extension to 90° SLR with brace unlocked
Weeks 9-12 weeks PT 1-2x/week HEP daily	Progress to full ROM Brace only when walking Return to normal ADLs	Maintain brace with ambulation Swimming OK at 12 weeks	Continue Phase 3 exercises Progress ROM as tolerated (active and passive) Closed chain resistance Progressive resisted quad strengthening until return to prior level of activity
Weeks 12+ PT 1-2x/week HEP daily	Maximize strength, flexibility, and endurance	Wean brace when quadriceps strength/control allows	Continue Phase 4 exercises Activity-specific exercises in effort to return patient to pre-injury level of activity